

Characterization of Alkaloids from The Leaves of *Psychotria Malayana* Jack of Lombok Island on The Basis of Gas Chromatography-Mass Spectroscopy

Surya Hadi^{1*}, Dina Asnawati¹, Vera Fitriya Ersalena¹, Ahmadun Azwari¹ and Khairul Pahmi Rahmawati¹


¹Study Program of Chemistry, Faculty of Mathematics and Natural Sciences, University of Mataram, Jl. Majapahit No. 62 Lombok, Indonesia.

Corresponding author: sur_hadi88@yahoo.com


GC-MS Chromatogram and Spectrum of Alkaloids from the Leaves Of *Psychotria Malayana* Jack from Lombok Island

Figure at injection temperature 100°C (compound 1-5)


A. Chromatogram of 1st injection at injection temperature 100°C in 35 minutes total running time



1. Chimonantine




Ion fragmentation of Meso-Chimonantine



Ion fragmentation of (+/-)- Chimonanthine


2. Calycanthine




Ion fragmentation of Calycanthine


B. Figure of 2nd injection at injection temperature 100°C in 45 minutes total running time

3. Hodgkinsine



Zoom in







Ion fragmentation of Hodgkinsine with various isomers

Figure at injection temperature 260°C (compound 6-7)

A. Figure of injection temperature 260°C in 20 minutes total running time




4. 2-ethyl-6-methylpyrazine



Ion fragmentation of 2-ethyl-6-methylpyrazine

5. 3-methyl-1,2,3,4-tetrahydro-gamma-carboline



Ion fragmentation of 3-methyl-1,2,3,4-tetrahydro-gamma-carboline